

d-Amphetamine Induced Changes in Social Interaction Patterns

EUCLID O. SMITH*¹ AND LARRY D. BYRD[†]

*Yerkes Regional Primate Research Center, *Departments of Anthropology and Biology and †Departments of Pharmacology and Psychology, Emory University, Atlanta, GA 30322*

Received 4 May 1984

SMITH, E. O. AND L. D. BYRD. *d*-Amphetamine induced changes in social interaction patterns. PHARMACOL BIOCHEM BEHAV 22(1) 135-139, 1985.—The behavioral effects of *d*-amphetamine were studied in a group of stump-tail macaques in a large outdoor enclosure. *d*-Amphetamine altered characteristic patterns of aggressive and affiliative behaviors in adult males that received the drug. Each monkey that received *d*-amphetamine increased its aggression toward non-adult animals in the group and decreased aggression toward adult members. In subjects for which genealogy was known, *d*-amphetamine increased aggression toward kin-related members of the group and decreased aggression toward non-kin monkeys. The effects of the drug on affiliative behaviors were less uniform and, therefore, less conclusive. Three subjects decreased affiliation and two increased affiliation toward non-adult monkeys. The results demonstrate that *d*-amphetamine can alter substantially the behavior of drug-treated members of a group and, in addition, that the drug can indirectly affect specific subsets of the group even though they did not receive the drug.

Stumptail macaques *d*-Amphetamine Aggressive behavior Objects of aggression

THE effects of *d*-amphetamine on behavior have been studied in a variety of species and with a number of different research protocols [7,13]. Although individual subjects studied in isolation have provided considerable knowledge about the behavioral effects of this and other drugs, recent research suggests that drug effects can also be influenced by the social environment of which the subjects are a part. For example, the effects of *d*-amphetamine can be influenced by the hierarchical or dominance position of a subject within a social group [19], and *d*-amphetamine can have qualitatively different effects on affiliative and aggressive behaviors in the same subjects in a group [20].

In a previous report, we characterized the effects of a range of doses of *d*-amphetamine on the rate of occurrence of aggressive and affiliative behaviors initiated by adult male monkeys living in a social group [20]. Over a wide range of doses, *d*-amphetamine decreased affiliative behavior in all drug-treated subjects, yet the drug increased aggressive behavior in the high- and low-ranking subjects. An extension of that work, the present report describes changes in social interactions, i.e., changes in the pattern or distribution of aggressive and affiliative behaviors toward specific members of the group, associated with *d*-amphetamine administration. The report identifies recipients of the behaviors and quantitates changes in the amount of aggressive or affiliative behavior directed toward recipients.

METHOD

Subjects

The experimental subjects were five adult male stump-tail

macaques (*Macaca arctoides*), weighing between 13.8 kg and 22.9 kg, and ranging in age from 5 to 11.5 years. The subjects were members of a heterogeneous group of 37 animals that included 5 adult males, 18 adult females, 3 sub-adult females, 3 juvenile males, 2 juvenile females and 6 immatures. The group was confined within a 28.4×32.7 m outdoor area with access to an environmentally-controlled, 4.4×12.2 m indoor area via two metal tunnels [21]. The positions of the five male subjects within the dominance hierarchy of the group ranged from the highest- to the lowest-ranking.

Procedures

All animals were restricted to the outdoor compound on a prescribed daily schedule, weather permitting, where each group member could be observed from a tower located 4.27 m above one side of the compound. Data characterizing the behavior of individual animals were collected and stored in a digital format using a microprocessor-based data collection device, the Datamyte 900, as described previously [18]. Subjects were observed and data were recorded during 15-minute test periods at preselected, post-injection times using the focal-animal sampling technique described by Altmann [1]. The focal-animal sampling technique permits the recording of all occurrences of various behaviors of an individual monkey during each sampling period. In this manner, a descriptive record can be obtained of an individual animal's behavior and of its interactions with other members of the group. In the present experiment, aggressive and affiliative behaviors initiated by drug-treated males toward other members of the group were studied. Affiliative behaviors

¹Requests for reprints should be addressed to Dr. Euclid O. Smith, Yerkes Regional Primate Research Center, Field Station, Emory University, 2409 Collins Hill Road, Lawrenceville, GA 30245.

TABLE 1
EFFECTS OF *d*-AMPHETAMINE ON AGGRESSIVE OR AFFILIATIVE BEHAVIORS INITIATED BY EACH OF FIVE ADULT MALES TOWARD ADULT VERSUS NON-ADULT MEMBERS OF THE GROUP

Subject	<i>d</i> -Amphetamine Dose		Non-adult	Adult	χ^2	<i>p</i>
Aggressive Behaviors						
M-13	0.56 mg/kg	Expected	45.9	118.1	0.50	≤ 0.50
		Observed	50.0 ↑	114.0 ↓		
M-10	0.03 mg/kg	Expected	17.1	39.3	1.37	≤ 0.30
		Observed	21.0 ↑	35.0 ↓		
M-06	0.03 mg/kg	Expected	32.3	53.6	16.12	≤ 0.001
		Observed	50.0 ↑	35.0 ↓		
M-24	0.03 mg/kg	Expected	3.1	35.5	123.88	≤ 0.001
		Observed	22.0 ↑	17.0 ↓		
M-18	0.01 mg/kg	Expected	26.9	70.1	21.40	≤ 0.001
		Observed	47.0 ↑	49.0 ↓		
Affiliative Behaviors						
M-13	0.56 mg/kg	Expected	6.3	30.7	7.58	≤ 0.01
		Observed	0.0 ↓	37.0 ↑		
M-10	0.30 mg/kg	Expected	25.2	58.8	2.94	≤ 0.10
		Observed	18.0 ↓	66.0 ↑		
M-06	0.30 mg/kg	Expected	15.2	110.5	3.27	≤ 0.10
		Observed	22.0 ↑	105.0 ↓		
M-24	0.30 mg/kg	Expected	6.0	54.6	6.80	≤ 0.01
		Observed	12.0 ↑	48.0 ↓		
M-18	0.30 mg/kg	Expected	9.7	78.3	1.57	≤ 0.25
		Observed	6.0 ↓	82.0 ↑		

were defined as any positive or affectionate behavior, or any behavior that was an attempt to safeguard another animal. Aggressive behaviors were defined as behaviors which could cause physical injury, signal the potential for harm, or result in preferential access to objects or events in the environment.

d-Amphetamine was dissolved in sterile normal saline (0.9%) and the resulting solution was injected intramuscularly in a volume of less than 1.0 ml. Sodium chloride solution (0.9%) served as a control (placebo) injection. Except for the highest dose (0.56 mg/kg) studied in monkey M-13 and the lowest dose (0.003 mg/kg) studied in monkey M-18, each dose was studied three times in each subject in an unsystematic order.

In order to inject the drug without alarming or exciting the animals, each of the five male subjects was trained to enter a small cubicle along one side of the outdoor compound [17] and to extend an arm through a circular opening according to procedures devised previously [3, 4, 15]. On a given day, each subject received either a drug injection, a saline (placebo) injection or no injection; however, only the experimental animal for that day received the drug. Persons responsible for data collection did not know whether saline or drug was administered. *d*-Amphetamine was administered two days per week, but a given animal was the experimental

or drug subject no more than once per week. Immediately after completing the injections, the subjects returned to the compound and data collection began. The initial observation period encompassed the entire first hour post-injection; thereafter, 15-minute observation periods began at 90, 150, 210 and 330 minutes post-injection.

RESULTS

Mean hourly rates of occurrence of affiliative and aggressive behaviors initiated by the male subjects following drug or saline administration were determined for each subject based on data obtained during the eight post-injection observation periods. Full dose-effect functions describing those data were reported previously [20]. In brief, the data on affiliative and aggressive behaviors initiated by the drug-treated animals showed overall monotonic decreases in affiliative behaviors by all five subjects as dose increased, and biphasic changes in aggressive behaviors in some subjects. To keep the mass of data at a manageable level for the present report, data presentation is limited to the dose of *d*-amphetamine that produced the maximum change in aggressive or affiliative behavior for each subject. *d*-Amphetamine can have differential effects depending on the dominance position of the drug-treated subject [20], and,

TABLE 2
EFFECTS OF *d*-AMPHETAMINE ON AGGRESSIVE AND AFFILIATIVE BEHAVIORS INITIATED BY TWO ADULT MALES TOWARD KIN VERSUS NON-KIN MEMBERS OF THE GROUP

Subject	<i>d</i> -Amphetamine Dose		Kin	Non-kin	χ^2	<i>p</i>
Aggressive Behaviors Initiated						
M-18	0.01 mg/kg	Expected	6.72	89.28	13.78	≤ 0.001
		Observed	16.00 ↑	80.00 ↓		
M-24	0.03 mg/kg	Expected	8.58	30.42	1.75	≤ 0.20
		Observed	12.00 ↑	27.00 ↓		
Affiliative Behaviors Initiated						
M-18	0.30 mg/kg	Expected	26.40	61.60	12.83	≤ 0.01
		Observed	11.00 ↓	77.00 ↑		
M-24	0.30 mg/kg	Expected	23.40	36.60	4.05	≤ 0.05
		Observed	31.00 ↑	29.00 ↓		

therefore, the dose that produced maximal effects differed among the subjects in the present experiment. Chi-square goodness-of-fit tests were used for all analyses; effects were considered significant if $p \leq 0.05$.

Age Analysis

The purpose of the present study was to describe effects of *d*-amphetamine on the directionality of affiliative and aggressive behaviors and to determine if the recipients of these behaviors represented special subsets of the larger social group. First, data were analyzed to determine if behavior was directed consistently toward non-adult as compared to adult members of the group and whether *d*-amphetamine altered the directionality of aggressive and affiliative behaviors. Adults were defined as individuals four years of age and older, and included those individuals that were reproductively mature. Non-adults included all monkeys less than four years of age. To apply the Chi-square test, the number of aggressive and affiliative interactions expected were determined based on the number of individuals in an age class and on frequencies of behavioral interactions following saline administration.

Table 1 summarizes data for the five males that received *d*-amphetamine. Although maximum increases in aggressive behaviors occurred at different doses for each of the five subjects, *d*-amphetamine uniformly increased aggressive behaviors toward non-adult members of the group and decreased aggressive behaviors toward adult monkeys in the group. The significance of the drug-induced changes was reflected in the results of the Chi-square tests, as shown in Table 1. The greatest redirection of aggressive behavior occurred in the three mid- and low-ranking animals, where values of $p \leq 0.001$ were obtained.

When the data were analyzed in a similar manner for maximum decreases in affiliative behaviors, a contrasting pattern resulted (Table 1). Three subjects decreased affiliative behaviors toward non-adult animals in the group and increased affiliative behaviors toward adults, and two subjects showed an opposite effect. The significance of the changes in recipients of affiliation yielded values of $p \leq 0.01$

for two of the five subjects when evaluated by Chi-square tests. However, none of the changes in affiliative behaviors was as large as the changes in recipients of aggressive behaviors.

Genealogical Analysis

To determine whether the recipients of aggressive or affiliative behaviors could be differentiated on the basis of blood relationship, data were examined for those drug-treated monkeys whose genealogy was known. Subjects for the genealogical analysis were monkeys M-18 and M-24, the two adult males that were born into the group and, therefore, the only two for which exact genealogy was known. Other group members were classified as kin-related or non-kin-related based on matrilineal records. Using rates of affiliative and aggressive behaviors under saline conditions and relative numbers of kin versus non-kin individuals in the group, expected rates of behavior for monkeys M-18 and M-24 were derived as shown in Table 2. Following the administration of *d*-amphetamine, both monkeys showed an increase in rate of aggressive behaviors toward kin-related animals and a decrease in aggression toward non-kin individuals.

d-Amphetamine also produced a significant change in recipients of affiliative behaviors following the administration of 0.30 mg/kg *d*-amphetamine. Statistically significant results were obtained for both subjects, indicating significant departures from expected values, but the effects were opposite in direction (Table 2). Monkey M-18 increased affiliative behaviors toward non-kin and monkey M-24 increased affiliative behaviors toward kin-related individuals.

DISCUSSION

The data presented here show that *d*-amphetamine can have pronounced effects on the expression of aggression and affiliation among animals living in a large, heterogeneous social group. When compared to the pattern of aggression and affiliation under baseline, non-drug conditions, significant changes in rates and in objects or recipients of the behaviors were found for both classes of behavior following administration of the drug.

Although previous research has revealed much regarding the effects of *d*-amphetamine on the behavior of individuals comprising various types of groupings [13], none has involved a group that yielded the type of analysis presented here. When the objects or recipients of aggressive behaviors were analyzed by age, for example, an unequivocal pattern based on age was revealed. Aggression directed toward young animals in the group increased for all five subjects following *d*-amphetamine administration. Moreover, the magnitude of the effect was inversely related to dominance position of the subjects in the group, with the highest-ranking male showing the smallest effect and the low-ranking males the largest effect. Differences in the statistical significance of the effects relative to dominance positions of the subjects indicated further that the magnitude of the effects might be due, in part, to the social position of each subject. These data are consistent with other reports showing that the behavioral effects of various drugs may be a function of the dominance position of the drug-treated subject. Wilson *et al.* [23] reported that *d*-amphetamine decreased food retrieval in high-ranking subjects and increased food retrieval in lower-ranking subjects. In other food competition tests, Lovell *et al.* [12], Grove *et al.* [8] and Bellarosa *et al.* [2] found increased food capture by subordinate animals when either the entire group or only the dominant animal was administered *d*-amphetamine. In other studies, Haber *et al.* [9,10] reported that *d*-amphetamine administration increased submissive behavior in low-ranking animals. Finally, Schlemmer and Davis [16] found rank-mediated changes in behavior following chronic *d*-amphetamine administration in small groups of stump-tail macaques.

Changes in the directionality of affiliative behaviors following drug administration were less uniform among the five subjects. Three of the subjects decreased and two increased the frequency of affiliative behaviors directed toward young animals in the group. A possible basis for this difference in effect among the five subjects was not readily apparent. The two subjects that increased affiliative behavior toward non-adult members of the group were in the low-middle positions of the dominance hierarchy, but there was little else to indicate that dominance position of the subject was a major determinant of the drug effect on affiliation. Indeed, we reported previously that *d*-amphetamine decreased affiliation uniformly and independently of dominance position in the group [20].

Of substantial interest in the present experiment were data showing that kin-related animals were more likely than non-kin-related animals to be targets of aggressive behaviors following *d*-amphetamine administration. Kinship groups have been shown to form strong interactional subunits for all

primate species where kinship relations among members of a social group were known [5, 6, 11, 14, 22]. This fact has been well documented by behavioral primatologists, but a detailed analysis of drug-induced behavioral changes in group-living animals with a known network of genealogical relations has not been reported. Although this phenomenon might normally be explained by greater spatial proximity to kin vs. non-kin, proximity did not account for the results when the data were corrected for a greater absolute number of non-kin animals. Furthermore, drug vs. non-drug comparisons were made against the saline control rate. The data were convincing in that both of the subjects with known genealogical histories exhibited the same pattern of increased aggression toward kin members of the group. That *d*-amphetamine had opposite effects on affiliation for the same two subjects indicates that there are probably other complex variables regulating the expression of affiliation in a socially-living primate.

The data reported here focus attention on the importance and value of studies to determine the effects of drugs on behaviors occurring naturally within a complex social group. The data show that the drug *d*-amphetamine not only increases the rate of aggressive behavior initiated by drug-treated members of the group, but that the drug-induced increase in aggressive behavior is distributed differentially among the various members of the group. The two characteristics that we have identified in this report as determinants of the expression of aggression are age and relationship of the recipients of the aggressive behavior. Younger, non-adult members of the group and kin-related members of the group received increased aggression following *d*-amphetamine administration. Finally, whereas previous reports have shown that *d*-amphetamine can affect differentially the rate of occurrence of aggressive and affiliative behaviors in group-living monkeys, the present report shows that *d*-amphetamine can significantly alter normal patterns of social relationships among adult and non-adult monkeys in a group. Disruption of patterns of social interactions can be as dramatic and as important as alterations in the rate of occurrence of naturally-occurring behaviors.

ACKNOWLEDGEMENTS

This research was supported by U.S. Public Health Service Grants DA-02128, RR-00165 and RR-00167 (Division of Research Resources, National Institutes of Health). We thank P. Peffer-Smith, S. Martin and P. Plant for assistance in preparation of the manuscript.

REFERENCES

1. Altmann, J. Observational study of behavior: sampling methods. *Behaviour* 49: 227-265, 1974.
2. Bellarosa, A., J. A. Bedford and M. C. Wilson. Sociopharmacology of *d*-amphetamine in *Macaca arctoides*. *Pharmacol Biochem Behav* 13: 221-228, 1980.
3. Byrd, L. D. Effects of *d*-amphetamine on schedule-controlled key pressing and drinking in the chimpanzee. *J Pharmacol Exp Ther* 185: 633-641, 1973.
4. Byrd, L. D. Introduction: chimpanzees as biomedical models. In: *Progress in Ape Research*, edited by G. H. Bourne. New York: Academic Press, 1977, pp. 161-165.
5. Chagnon, N. A. Genealogy, solidarity, and relatedness: limits to local group size and patterns of fissioning in an expanding population. *Yrbk Phys Anthropol* 19: 95-110, 1975.
6. Chepko-Sade, B. D. and D. S. Sade. Patterns of group splitting within matrilineal kinship groups: a study of social group structure in *Macaca mulatta* (Cercopithecidae: Primates). *Behav Ecol Sociobiol* 5: 67-86, 1979.
7. Costa, E. and S. Garattini, editors. *International Symposium on Amphetamines and Related Compounds*. Proceedings of the Mario Negri Institute for Pharmacological Research, Milan, Italy. New York: Raven Press, 1970.

8. Grove, L., M. Wilson and J. Bedford. Effects of diazepam and d-amphetamine on food competition in rhesus monkey. *Pharmacologist* **19**: 228, 1977.
9. Haber, S. N., P. R. Barchas and J. D. Barchas. Effects of amphetamine on social behaviors of rhesus macaques: an animal model of paranoia. In: *Animal Models in Psychiatry and Neurology*, edited by I. Hanin and E. Usdin. Elmsford, NY: Pergamon Press, 1977, pp. 107-115.
10. Haber, S. N., P. A. Berger and P. R. Barchas. The effects of amphetamine on agonistic behaviors in nonhuman primates. In: *Catecholamines: Basic and Clinical Frontiers*, edited by E. Usdin, I. J. Kopin and J. Barchas. Elmsford, NY: Pergamon Press, 1979, pp. 1702-1704.
11. Koyama, N. Changes in dominance rank and division of a wild monkey troop in Arashiyama. *Primates* **11**: 335-390, 1970.
12. Lovell, D. K., J. A. Bedford, L. Grove and M. C. Wilson. Effects of d-amphetamine and diazepam on paired and grouped primate food competition. *Pharmacol Biochem Behav* **13**: 177-181, 1980.
13. Miczek, K. A. (ed) *Ethopharmacology: Primate Models of Neuropsychiatric Disorders*. New York: Alan R. Liss, 1983.
14. Nash, L. T. Troop fission in free-ranging baboons in the Gombe Stream National Park, Tanzania. *Am J Phys Anthropol* **44**: 63-77, 1976.
15. Peffer-Smith, P. G., E. O. Smith and L. D. Byrd. Effects of d-amphetamine on self-aggression and posturing in stump-tail macaques. *J Exp Anal Behav* **40**: 313-320, 1983.
16. Schlemmer, R. F., Jr. and J. M. Davis. Evidence for dopamine mediation of submissive gestures in the stump-tail macaque monkey. *Pharmacol Biochem Behav* **14**: 95-102, 1981.
17. Smith, E. O. Device for capture and restraint of monkeys. *Lab Anim Sci* **31**: 305-306, 1981.
18. Smith, E. O. and M. L. Begeman. BORES: Behavior observation recording and editing system. *Behav Res Meth Instrum* **12**: 1-7, 1980.
19. Smith, E. O. and L. D. Byrd. Studying the behavioral effects of drugs in group-living nonhuman primates. In: *Ethopharmacology: Primate Models of Neuropsychiatric Disorders*, edited by K. A. Miczek. New York: Alan R. Liss, 1983, pp. 1-31.
20. Smith, E. O. and L. D. Byrd. Contrasting effects of d-amphetamine on affiliation and aggression in monkeys. *Pharmacol Biochem Behav* **20**: 255-260, 1984.
21. Smith, E. O. and P. G. Peffer-Smith. Adult male-immature interactions in captive stump-tail macaques (*Macaca arctoides*). In: *Primate Paternalism: An Evolutionary and Comparative View of Male Investment*, edited by D. M. Taub. New York: Van Nostrand Reinhold, 1984, pp. 88-112.
22. van Lawick-Goodall, J. Mother-offspring relationships in free-ranging chimpanzees. In: *Primate Ethology*, edited by D. Morris. Chicago: Aldine Publ. Co., 1967, pp. 287-346.
23. Wilson, M. C., L. Bailey and J. A. Bedford. Effects of subacute administration of amphetamine on food competition in primates. In: *Ethopharmacology: Primate Models of Neuropsychiatric Disorders*, edited by K. A. Miczek. New York: Alan R. Liss, 1983, pp. 293-305.